

The Decomposition of Organic Compounds at High Temperatures and Pressures

A DISSERTATION

SUBMITTED TO THE BOARD OF UNIVERSITY STUDIES OF THE
JOHNS HOPKINS UNIVERSITY IN CONFORMITY WITH THE
REQUIREMENTS FOR THE DEGREE OF DOCTOR
OF PHILOSOPHY

BY
LEE ROY HERNDON
Baltimore, 1928

EASTON, PA.:
MACK PRINTING COMPANY

1928

The Decomposition of Organic Compounds at High Temperatures and Pressures

A DISSERTATION

SUBMITTED TO THE BOARD OF UNIVERSITY STUDIES OF THE
JOHNS HOPKINS UNIVERSITY IN CONFORMITY WITH THE
REQUIREMENTS FOR THE DEGREE OF DOCTOR
OF PHILOSOPHY

BY
LEE ROY HERNDON
Baltimore, 1928

EASTON, PA.:
MACK PRINTING COMPANY
1928



TABLE OF CONTENTS

Introduction.....	5
Apparatus and Procedure.....	5
Results.....	6
Summary.....	12
Biography.....	13

ACKNOWLEDGMENT

The author takes this opportunity to express his appreciation to Professor E. Emmet Reid at whose suggestion and under whose capable direction the investigations have been carried out, for his classroom instruction, as well as many practical suggestions in the laboratory.

Thanks are also due Dr. F. R. Bichowsky for securing part of the apparatus and for valuable advice in construction of the apparatus, as well as for laboratory and classroom instruction.

The author wishes to express his appreciation to Professors Frazer and Patrick, and Associate Professor Rice, and Doctors Andrews, Urey and Thornton for their valuable instruction in lectures and laboratory.

THE DECOMPOSITION OF ORGANIC COMPOUNDS AT HIGH TEMPERATURES AND PRESSURES¹

Introduction

The purpose of this investigation was the determination of the effect of high temperatures upon organic compounds at constant volume.

Numerous experiments have been carried out on decomposition of organic compounds at high temperatures by various methods, such as passing the vapors through hot tubes in the presence of catalysts or by electrical discharge, but, for the most part, these have been carried out under atmospheric pressure. Sealed tubes and autoclaves have been used but these are not designed to withstand extremes of temperature and pressure. Very few experiments have been made at temperatures above 250° and at pressures greater than a few hundred atmospheres. It seemed desirable to study the decomposition of pure organic compounds when heated at constant volume and the consequent pressure at temperatures from 250–525°.²

Apparatus and Procedure

The apparatus used in these experiments was designed to withstand a pressure of 20,000 atmospheres at 400°, at which temperature the tensile strength of the chrom-vanadium³ steel used is at a maximum. The design was such that at the weakest place in the structure the wall was three-fourths of an inch in thickness. The bomb which was used was of the same design as that used by Morey⁴ except that the chamber was made longer so as to contain 100 cc. A thermocouple well was bored down the center of the plunger to within one-half inch of the reaction chamber. The bomb was heated in an electric furnace which was controlled by means of an automatic regulator⁵ to which the thermocouple was connected, and in series with three lamp-banks, two of which were connected to the regulator through relays. By this means it was possible to maintain any given temperature between 250 and 525°, within $\pm 2^\circ$ for any desired length of time.

The threads were dusted with graphite, which was used as a lubricant. The bomb was then sealed with a copper gasket and made tight by means of a long handled, double ratchet wrench. The heating was usually for two hours for the initial run, the

¹ Presented at the Richmond meeting of the American Chemical Society, April, 1927.

² This was made possible by the use of a type of bomb developed in the Geophysics Laboratory and brought to our attention by Dr. F. R. Bichowsky.

³ Procured from the Midvale Steel Co., Philadelphia, Pa.

⁴ Morey, *J. Am. Chem. Soc.*, **36**, 215 (1914).

⁵ Leeds and Northrup Co., Philadelphia, Pa.

time being lengthened if necessary to effect decomposition. At the desired time the bomb was removed from the furnace and allowed to cool spontaneously to room temperature. It was then set in the holding device, and the wrench applied very slowly until the gaseous products could be heard to escape, then the wrench was quickly removed and the bomb transferred to the water-bath and gas collecting apparatus. This was an inverted 14-liter bottle filled with water. Through the stopper passed a funnel and a tube to discharge the water displaced by the gas. After the gas had escaped the bomb was taken from the bath, dried, opened and the liquid products were removed by means of a pipet. If there were two layers, these were weighed separately. The gaseous products were measured and analyzed and the liquid products subjected to distillation and the products identified as far as possible.

The bombs withstood all pressures to which they were subjected except in one case. In one run with ethylene glycol at 400° the bomb burst. A slit about three inches long opened in the side. The furnace was wrecked but little other damage was done as there were no flying pieces of metal. The bursting was probably due to the thermal expansion of the liquid glycol due to the bomb being too nearly filled with the charge.

Results

Alcohols.—The aliphatic alcohols, methyl-, ethyl-, propyl-, isopropyl-, *n*-butyl-, isobutyl-, *sec*-.butyl-, *tert*-.butyl- and *n*-amyl-; the aromatic, benzyl- and phenylethyl; and the terpene, fenchyl alcohol have been decomposed by heating at temperatures above 400° for two hours. Temperatures were chosen such that decomposition was not quite complete in this time. All of the above mentioned alcohols yielded a great variety of products. There seem to be two simultaneous reactions: (1) dehydration and subsequent polymerization of the hydrocarbon residues, and (2) dehydrogenation followed by the splitting off of carbon monoxide and then polymerization. Three distinct types of compounds were derived from these decompositions: (1) gaseous, (2) water-insoluble portion, (3) water and water-soluble portion.

The gaseous products were analyzed in a number of cases, the results of which are given in the following table.

TABLE I
PERCENTAGE COMPOSITION OF THE GASEOUS PRODUCTS FROM ALCOHOLS



Alcohol	CO ₂	CH ₄	CH ₂ =CH ₂	CO	C ₂ H ₆	H ₂
Methyl	8.1	41.8	0.8	3.1	..	39.4
Ethyl	10.2	31.8	8.0	7.6	36.6	..
<i>n</i> -Propyl	9.8	36.8	2.0	21.6	..	20.0
Isopropyl	8.6	..	18.4	2.8	51.2	12.6
Benzyl	14.2	4.2	0.6	64.6	..	11.0
Phenylethyl	10.4	5.8	..	44.6	..	33.0
Fenchyl	10.0	32.2	2.4	51.0	..	..

Methyl alcohol gave exceptional results in that no liquid hydrocarbon or insoluble portion was obtained, the products of decomposition being gaseous with the exception of water. Peytral⁶ and Ipatiev⁷ state that


⁶ Peytral, *Compt. rend.*, **165**, 703 (1917).

⁷ Ipatiev, *Ber.*, **34**, 3579 (1901); **35**, 1047 (1902).

the products are all gaseous, the decompositions taking place according to the following reactions



In the bomb the above reactions appear to be accompanied by hydrogenation. That reaction (2) takes place is proved by the presence of carbon monoxide in the gaseous product; however, the dioxide is present to a greater extent than the monoxide. This may be accounted for by the reaction:



which Briner and Wroczynski⁸ found to take place at 320–330° and 400 atmospheres.

Ethyl alcohol showed dehydrogenation, dehydration and polymerization of the hydrocarbon residue to compounds of wide boiling range, as shown in Table II, which also shows the density and bromine absorption for each cut.

TABLE II

FRACTIONAL DISTILLATION OF 591 G. OF UPPER LAYER FROM 30 RUNS OF 59 G. EACH OF ETHYL ALCOHOL AT 425° FOR TWO HOURS

Fraction	T, °C.	G.	Density	Br abs., g. per cc.	Frac- tion	T, °C.	G.	Density	Br abs., g. per cc.
1	18–30	22	0.6252	0.1122	10	138–160	31	0.7751	0.2019
2	30–40	24	.6380	.1047	11	160–166	38	.7936	.2188
3	40–50	32	.6830	.1327	12	166–195	43	.8134	.1927
4	50–63	107	.7034	.1421	13 ^a	90–110	47	.8430	.1627
5	63–80	22	.7120	.2468	14 ^a	110–130	28	.8509	.1208
6	80–98	32	.7252	.1701	15 ^a	130–150	15	.8526	.1458
7	98–106	28	.7430	.2262	16 ^a	150–170	15	.8591	.1645
8	106–119	28	.7552	.2337	17 ^a	170–200	13	.9243	.1533
9	119–138	51	.7647	.2075	18 ^a	200–270	15	.9710	.0195

^a At 15 mm.

Unsaturation was also evident in the gaseous product. This unsaturation cannot be entirely credited to the lack of hydrogen in the decomposition products because the gaseous portion from isopropyl alcohol was even more unsaturated than that from ethyl alcohol and still contained about 12% hydrogen.


Sec.-butyl alcohol was exceptional in so far as no gaseous product was obtained.

Tert.-butyl alcohol was almost completely decomposed when heated at 400° for two hours, yielding nearly the theoretical quantity of water, showing that the primary reaction is dehydration. This was the lowest temperature at which any of the alcohols underwent complete decomposition.



⁸ Briner and Wroczynski, *Compt. rend.*, **150**, 1324 (1910).

Benzyl alcohol yielded carbon monoxide as the chief gaseous product. The toluene formed must have resulted from hydrogenation, since both water and hydrogen were present in the final products of the decomposition.

Hydrocarbons.—Normal heptane, the only aliphatic hydrocarbon subjected to 400° for two hours, underwent slight decomposition into simpler products as well as some of more complex character. The simpler products were gaseous, while the most complex were of the character of lubricating oil. The mechanism given by Thorpe and Young⁹ for butane is



but Pease¹⁰ has shown it to be



In the bomb the normal saturated hydrocarbon yields simpler saturated hydrocarbons together with unsaturated hydrocarbons, which under the influence of heat and pressure polymerize to give higher hydrocarbons. Each fraction contained some unsaturates, since each took up an appreciable quantity of bromine.

Benzene is quite stable at 400° and was only slightly decomposed when heated at 500° for two hours. More complete decomposition was effected by heating at 525° for forty-eight hours, but even then the decomposition was incomplete, 27% of the benzene being recovered. With the exception of the hydrogen the decomposition products were solids. Diphenyl, *m*-diphenyl- and *p*-diphenylbenzenes were formed in quantities sufficient for identification, while other products of higher boiling point did not yield readily to the action of solvents or the formation of derivatives.

Toluene decomposed almost completely when heated at 525° for ten hours, with the deposition of some carbon. Distillation of the resulting product indicates the formation of xylenes and a number of isomers with the empirical formula $\text{C}_{14}\text{H}_{14}$, which boil over the range of 278–300°.

Other aromatic hydrocarbons with side chains, such as ethylbenzene, *isopropylbenzene*, *tert.*-butylbenzene and unsymmetrical diphenylethane behave much as does toluene.

Acids.—Acetic acid at temperatures above 400° decomposes into gaseous products rather than yielding liquid products of varied boiling points, as is shown by the distillation range of the product.

Adipic acid at 450° for three hours breaks down completely into liquid and gaseous products with the deposition of carbon. The liquid portion was in two layers, the upper being a dark brown oil, while the lower layer was colorless and mostly water.

Benzoic acid heated at 400° for two hours yielded a thick, black sub-

⁹ Thorpe and Young, *Proc. Roy. Soc.*, **21**, 184 (1873).

¹⁰ Pease, *J. Am. Chem. Soc.*, **50**, 1779 (1928).

stance which was chiefly undecomposed benzoic acid with a little carbon and some gas as decomposition products.

Citric acid decomposes at 400° into two liquid layers, a large quantity of gas and a little carbon. The upper layer was composed of hydrocarbons, while the lower layer contained acetone, acetic acid and water.

Stearic acid when heated at 400° for two hours gave a pale yellow, soft solid which appeared to be the unchanged acid.

Aldehydes.—Acetaldehyde at 335° for two hours gave two liquid layers of almost equal volume and two liters of gas. The upper layer gave a wide variety of products, as indicated by the boiling range of 55 – 360° . The lower layer distilled over a narrow range, 79 – 100° , the major portion within the last 5° . The gaseous product contained carbon dioxide, carbon monoxide, unsaturated hydrocarbons and methane.

Benzaldehyde at 400° for two hours gives benzene as the principal product as well as other higher boiling products.

Chloroform.—Chloroform decomposed at 425° yielding chlorine, and possibly hydrogen chloride, with two liquid layers and a pale yellow solid which had the correct melting point for hexachloro-ethane. Work was discontinued on this product because of the corrosive effect of the products of decomposition on the bomb.

Ester.—Ethyl acetate above 400° for two hours decomposes into two liquid layers, a little gas and a small amount of carbon. The upper layer, consisting mostly of hydrocarbons, had a very wide boiling range, indicating a large number of compounds. The lower layer was chiefly water with some acetic acid formed by hydrolysis and a soluble ferrous salt, possibly the acetate.

Ether.—Diethyl ether decomposes when heated at 400° for two hours yielding acetaldehyde, considerable gas and a liquid distilling over the range of 22 – 48° .

Glycols.—Ethylene glycol decomposes into two liquid layers and some gas when heated at temperatures considerably below 400° .

Diethylene glycol undergoes almost complete decomposition at 400° for two hours. The products of decomposition consist of some gas and two immiscible liquids. The liquids show a very marked change in character from the original. In addition to having distillation ranges lower than the boiling point of pure diethylene glycol, the lower layer is chiefly water, as is the case with the lower layers of the alcohols, while the upper layer is composed of hydrocarbons of a wide boiling range.

Acetone.—Acetone at 400° for two hours decomposes into two layers, the lower of which is chiefly water and the upper layer of an oily consistency.

Benzophenone.—Benzophenone appears to be quite stable at 400° for two hours.

Phenol.—Phenol is stable at 400° and decomposes only slightly when heated at 525° for two and one-half hours.

Quinol.—Quinol undergoes some decomposition at 400° in two hours. The product from one experiment was not of sufficient quantity to justify attempts at identification.

Sucrose.—Cane sugar gave a black, porous mass when heated at 425° for two hours.

TABLE III
SUMMARY OF RESULTS

Cpd.	Temp., °C.	Time, hours	Charge, g.	Product Upper	Lower	Gas, liters	Dist. range, °C.	Cpds. identified and remarks
Methyl alcohol	400	2	60	39	..	10	64–100	{ H ₂ O, CH ₃ OH, CO ₂ , CO, CH ₂ , H ₂
	435	2	50	20	..	Much		
	450	2	60	20	..	15		
	475	2	60	48	..	Much	64–99	
Ethyl alcohol	425	2	59	17f ^a	15	14.5	Upper, 18–270;	{ CH ₃ COOH, CH ₃ CHO H ₂ O, CO, CO ₂ , CH ₄ , C ₂ H ₄ , C ₂ H ₆
	460	2	59	9f	15	19	lower, 30–102	
	425 ^b	2	59	19.7f	16.7	14.5		
Propyl alcohol	400	2	40	40	..	...		One layer
	415	2	40	29	..	Little	27–115	
	425	2	40	19f	7	Much	27–105	{ C ₂ H ₅ CHO, H ₂ O, CO ₂ , CO, H ₂ , C ₂ H ₄ , CH ₄
	425	2	48	30f	7	Much	35–165	
	425	2	60	12f	7	11.5		
Isopropyl alcohol	400	2	42	42	..	...	79–82	
	425	2	42	23	..	Little	57–85	
	425	2	42	3f	29	...		{ CH ₃ COCH ₃ , H ₂ O, CO ₂ , C ₂ H ₄ , CO, C ₂ H ₆ , H ₂
	425	8	42	11f	15r	...		
	435	8	42	7f	117	...		
	465	.	52	16f	16	Much	56–100	
	435	2	60	26f	26	3	35–112	
Butyl alcohol	400	2	40.5	40	..	...	65–119	Colorless liquid, H ₂ O
	435	2	40.5	21f	4	Much	52–200	
Isobutyl alcohol	400	2	60	42	..	Little	40–112	
	400	4	60	11f	1	Much		
	400	16	60	8f	3	Much		
	425	2	60	7f	2	Much		
	450	2	60	8f	2	Much		
	475	2	60	3f	2	18		
Sec.-butyl alcohol	400	2	41	41	..	None	23–102	C ₂ H ₅ OH
	425	2	41	38f	3	...		
Tert.-butyl alcohol	400	2	39	20f	11	...	26–200	
	300	2	39	26	..	...	80–98	f liquid
n-Amyl alcohol	425	2	60	35f	10	5	30–215	H ₂ O
	450	2	60	38f	5	Little	35–215	
	450	16	15	7	3	18		
Benzyl alcohol	425	2	50	38	5	3	78–305	{ C ₆ H ₅ , C ₆ H ₅ CH ₃ , BzOH, CO ₂ , CO, C ₂ H ₄ , CH ₄ , H ₂
	425	2	50	43	5	Little	83–100	

TABLE III (Continued)

Cpd.	Temp., °C.	Time, hours	Charge, g.	Product		Gas, liters	Dist. range, °C.	Cpds. identified and remarks
				Upper	Lower			
Phenylethyl alcohol	425	2	50	42	6	1.5	60-330	$\left\{ \begin{array}{l} \text{CH}_3\text{OH}, \quad \text{C}_6\text{H}_5\text{CH}_3 \\ \text{CO}_2, \text{CH}_4, \text{H}_2, \text{CO}, \\ \text{H}_2\text{O} \end{array} \right.$
	425	2	50	43	6	Little		
Fenchyl alcohol	435	2	75	57.3	1	2.5	30-335	$\text{H}_2\text{O}, \text{CO}_2, \text{C}_2\text{H}_4, \text{CO},$ CH_4
Acetic acid	350	2	79	49	..	...	114-118	
	412	2	79	41	..	3.7	110-118	
Adipic acid	450	3	60	3	15	Much		
Benzoic acid	400	2	15	12	..	Little		
Citric acid	400	2	50	10	8	Much		$\text{H}_2\text{O}, \quad \text{CH}_3\text{COCH}_3,$
	250	2	50	14	10	Much		CH_3COOH
Stearic acid	400	2	15	15	..	Little		Yellow solid
Acetaldehyde	425	2	60	8	12	Much		
	325	2	60	20	17	2	55-360	$\text{CO}_2, \text{C}_2\text{H}_4, \text{CO}, \text{CH}_4$
Benzaldehyde	400	2	52	33	..	Much	81-297	$\text{C}_6\text{H}_6 \quad \text{C}_6\text{H}_5\text{COOH}$
	400	2	52	36	..	Much	81-297	
<i>n</i> -Heptane	400	2	52	29	..	5	27-295	
Benzene	400	2	66	55	..	0.5	68- 80	f liquid
	400	24	66	66	..	...	75- 80	f liquid
	450	24	66	50	..	...	70- 80	f liquid
	500	2	66	63	..	Little	69- 82	f liquid
	600	.	66	17	..	...	250-280	Solid
	500	18	66	60	..	5.5	70-310	Solid
	500	24	66	66	..	...	80-254	f liquid, diphenyl
	525	4	66	65	..	...	80-500	f liquid
	525	24	66	59	..	Little	80-260	Liq., solid diphenyl
	525	48	66	50	..	Much	80-500	$\left\{ \begin{array}{l} \text{H}_2, \quad \text{C}_6\text{H}_5\text{C}_6\text{H}_5, \quad m\cdot \\ \text{C}_6\text{H}_4(\text{C}_6\text{H}_5)_2, \quad p\cdot \\ \text{C}_6\text{H}_4(\text{C}_6\text{H}_5)_2 \end{array} \right.$
	525	48	66	55	..	Much	80-500	
	525	48	66	58	..	Much	80-500	
Toluene	400	2	43	43	..	...	108-110	
	500	2	43	43	..	...	97-110	
	525	10	57	54	..	...	100-300	C_6H_6
	525	48	57	35	..	...		
Ethylbenzene	400	2	44	43	..	...	130-138	
	400	8	58	54	..	...	130-140	
	450	2	58	51	..	...	125-145	
	450	2	66	66	..	...	124-140	
	450	10	66	66	..	Little	108-140	
	450	24	58	27	..	Much		Solid, 14 g.
	500	2	66	41	..	...	97-140	$\text{C}_6\text{H}_6,$
	500	2	66	66	..	Little	105-305	$\text{C}_6\text{H}_5\text{CH}_3$
Isopropyl benzene	450	3	57	50	..	Little	140-200	
<i>Tert.</i> -butyl- benzene	450	2	59	51	..	Little	105-175	$\text{C}_6\text{H}_5\text{CH}_3$
Diphenyl- ethane	425	16	55	34	..	2	80-360	

TABLE III (Concluded)

Cpd.	Temp., °C.	Time, hours	Charge, g.	Product		Gas, liters	Dist. range, °C.	Cpds. identified and remarks
				Upper	Lower			
Chloroform	425	2	112	35	23	...	100-109	Cl ₂ , HCl, C ₂ Cl ₆
Ethyl acetate	400	2	42	19	11	Little	39-178	CH ₃ COCH ₃ , CH ₃ - COOH, H ₂ O, CH ₃ - CHO
	425	2	42	18	9	Little	33-200	
Ether	400	2	36	18	..	Much	22-48	CH ₃ CHO
Glycol	400	2	64	15	19	13.6	68-101	
	350	4	84	2	82	1.5	67-205	
Diglycol	400	2	56	10	14	Much	50- 90	
	400	2	56	10	24	5	65-101	C ₂ H ₅ OH, H ₂ O
	400	22	56	8	18	Much		
Acetone	400	2	26	26	..	...	56- 60	
	400	2	39	30	5	Little	53-170	H ₂ O
Benzophenone	400	2	37	36	..	...	293-310	
Phenol	400	2	50	50	..	...		
	500	2.5	50	50	..	...	80-187	
Quinol	400	2	44	38	..	Little	98-280	
Sucrose	425	2	15	15	..	...		

^a The figures for the third run with ethyl alcohol are the average of those obtained in 30 runs under the same conditions.

^b f = fluorescent.

Summary

A study has been made of the decomposition of organic compounds at constant volume, constant temperature (between the limits of 300-525°), at consequent pressure and for periods of time ranging from two to forty-eight hours, the temperature and duration of heating depending upon the stability of the compound under investigation.

The following substances have been heated: methyl-, ethyl-, *n*-propyl-, isopropyl-, *n*-butyl-, isobutyl-, *sec*-.butyl-, *tert*-.butyl-, *n*-amyl, benzyl-, phenylethyl and fenchyl alcohols; acetic, adipic, benzoic, citric and stearic acids; acetaldehyde and benzaldehyde; *n*-heptane; benzene, toluene, ethylbenzene, isopropylbenzene, *tert*-.butylbenzene; unsymmetrical diphenylethane; chloroform; ethyl acetate; diethyl ether; ethylene and diethylene glycols; acetone and benzophenone; phenol; quinol; and sucrose.

The products of decomposition have been separated and identified as far as possible.

BIOGRAPHY

Lee Roy Herndon, the author of this dissertation, was born at Welchburg, Kentucky, October 31, 1897. He received his early education in the public schools of Jackson County, Kentucky, graduating from Junior High School at Annville Institute, Annville, Kentucky, in 1917. He entered the preparatory department of Maryville College, Maryville, Tennessee in September, 1917, from which department he graduated in 1919. His college work was also done at Maryville College, from which he received the A.B. degree in 1922. The year 1922-1923, he taught science in the Central High School of Lewisburg, Tennessee. The summers of 1923 and 1924 were spent in the University of Chicago, Chicago, Illinois. The scholastic years of 1923-1925, he was professor of Chemistry in Maryville College. In September 1925, he entered Johns Hopkins University where he has been engaged in the work discribed in the foregoing pages.

